



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Subjects.	Ether.	Orange.	Romarin.	Wintergreen.	Menth.	Vanillin.
No. 1 $\text{Æ}$	27 0.5	0.05	0.05	0.005	0.0005	0.0006
No. 2 "	22 1.	0.05	0.55	0.005	0.01	0.0005
No. 3 "	32 1.	0.05	0.05	0.005	0.001	0.00007
No. " "	42 1.	1 to 3	0.05	0.05	0.005	0.001
No. " "	40 4.	1 to 3	0.8	0.4	0.01	0.005

The different stages of the experiment may be defined as 1, smelling nothing; 2d, smelling something that it is not possible to define, and 3, ability to name the odor correctly. The figures found agree quite closely with those determined by Valentin many years ago.

*On the nerve supply of the sense of taste*, John Ferguson, M. A., M. D.—Medical News, 1890 Vol. LVII. 395.

It is quite generally admitted that the lingual branch of the fifth nerve and the gustatory branches of the glossopharyngeal carry the nerve fibres of taste to the tongue and palate. Are these the real supply to the parts of taste, or do they simply carry nerve fibres of taste to the tongue and palate? The author has had an excellent opportunity to observe a case in which there was a total loss of taste on the left side of the tongue, even to the tip. An autopsy was made which proved that the nerve supply of taste for the top and anterior part of the sides of the tongue comes from the fifth nerve and enter the superior maxillary division of the same nerve. The course then must be from the superior maxillary nerve into the spheno-palatine ganglion, thence by the vidian through the vidian canal to the gangliform enlargement of the facial, along this to the chorda tympani, through the chorda tympani into the lingual, a branch of the inferior maxillary of the fifth.

The conclusion is also reached that the vidian is not a motor root passing from the facial to the spheno-palatine ganglion, but a sensory nerve of the special sense of taste from the spheno-palatine ganglion of the second division of the fifth nerve to the seventh or facial; also nerve fibres of taste for the back of the tongue, fauces and soft palate cannot be carried by the chorda tympani. Proof is also adduced that the root of the glossopharyngeal nerve does not contain any fibres of the special sense of taste.

The route then for the sense of taste, so far as the glossopharyngeal is concerned, would be from the root of the fifth through its third division to the otic ganglion, from this by the small petrosal to the ganglion of the seventh, thence to the tympanic plexus, again by the tympanic branch to the petrous ganglion of the glossopharyngeal and by this latter to the back of the tongue, fauces and palate.

In a criticism on the above article in the Medical News, Vol. LVII. p. 464, by Dr. Chas. L. Dana, he asserts that while the researches noticed seem to prove that loss of taste is due to a disease of the vidian nerve, they do not prove so conclusively that the glossopharyngeal nerve has no gustatory functions.

*Sur la physiologie comparée de l'olfaction*, M. Raphael Dubois.—Comptes Rendus, Vol. CXI. 66.

The olfactory organs of mollusks have been studied by numerous experimenters, but so far little has been learned of the mechanism of the organs or of their mode of acting. It is a fact that many odors excite the organ of smell of *Helix Pomatia*, a mollusk well adapted to experiments on the special senses. The large tentacles are more sensitive than other parts, while that of the small tentacles, though considerable, is notably less than that of the large ones. As far as the rest of the external covering of the mollusk is concerned, it is excited by only a few odoriferous substances. There is greater sensibility at the extremity of the large tentacles, though it is apparent throughout the